

Date of publication: 18 January 2016

Implementation: To be determined by each Service

## **Change Notification UK National Blood Services No. 04 2016**

This change applies to All the Donor Selection Guidelines

### **Table of Immunizations**

Please add this new Appendix 2

<b>Diseases protected against</b>	<b>Comments and example trade names of adult preparations</b>	
Anthrax	Rarely given available only through Public Health England.	<u>Non-Live</u>
Cholera	Oral vaccine. Dukoral®	<u>Non-Live</u>
Diphtheria	Would usually be given as Diphtheria, Tetanus, Pertussis often with Polio in older children and adults, Boostrix-IPV®, Revaxis®, Repevax®	<u>Non-Live</u>
HiB	Haemophilus influenza, Menitorex®	<u>Non-Live</u>
Hepatitis A	May be combined with typhoid or hepatitis B. Avaxim®, Epaxal®, Ambirix®, Twinrix®, Hapatyrix®, Viatim®	<u>Non-Live</u>
Hepatitis B	May be combined with hepatitis A If unexposed and more than 7 days from last immunisation, accept. See: <u>Hepatitis B – Immunisation</u> Engerix®, Fendrix®, HBvaxPRO®, Ambirix®, Twinrix®	<u>Non-Live</u>
HPV	Usually given to teenage girls, Cervarix®, Gardasil®	<u>Non-Live</u>
<b>Influenza-intra-nasal</b>	<b>Fluenz® to age 18</b>	<b><u>Live</u></b>
Influenza	Injection-annual flu jab. Agrippal®, Enzira®, Fluarix®, Fluvirin®, Imuvac®, Influvac®, Intanza® Optaflu®, Pandemrix®	<u>Non-Live</u>
Japanese Encephalitis	Travel. Ixiaro®	<u>Non-Live</u>
<b>Measles, Mumps, Rubella</b>	<b>MMR vaccines. M-M-RvaxPro®, Priorix®</b>	<b><u>Live</u></b>
Meningitis	Meningococcal group C, Meningococcal group B and the Quadrivalent ACWY immunisation may also be given in combination with HiB. Meningitec®, Menjugate Kit®, NeisVac-C®, Bexsero®, Menveo®, Nimenrix®	<u>Non-Live</u>

**Continued**

Pertussis	Usually pregnant women, Boostrix-IPV®	<u>Non-Live</u>
Pneumococcal disease	Given to people with specific risks e.g Asplenic. And over 65s Pneumovax II®	<u>Non-Live</u>
Polio	Would usually be given Diphtheria, Tetanus and Polio (injection). Boostrix-IPV®, Revaxis®, Repevax®	<u>Non-Live</u>
Rabies	Travel or post-bite. Rabipur®	<u>Non-Live</u>
<b>Shingles</b>	<b>Routine immunisation for older people. Zostavax®</b>	<u>Live</u>
<b>Smallpox</b>	<b>Rarely given, see: <u>Smallpox Immunization</u></b>	<u>Live</u>
Tetanus	Would usually be given Diphtheria, Tetanus and Polio. Revaxis®	<u>Non-Live</u>
Tick-borne encephalitis	Tico-vac®	<u>Non-Live</u>
<b>Tuberculosis</b>	<b>BCG vaccine</b>	<u>Live</u>
Typhoid - injected	Typherix®, Typhim Vi®	<u>Non-Live</u>
<b>Typhoid - Oral</b>	<b>Vivotif®</b>	<u>Live</u>
<b>Varicella (chickenpox)</b>	<b>Mainly healthcare workers. Varilrix®, Varivax®,</b>	<u>Live</u>
<b>Yellow Fever</b>	<b>Travel. Stamaril®</b>	<u>Live</u>

*Sheila MacLennan*

Dr Sheila MacLennan  
Professional Director - Joint UKBTS Professional Advisory Committee  
☎ Direct Dial: (0113) 820 8638    ✉ [sheila.maclennan@nhsbt.nhs.uk](mailto:sheila.maclennan@nhsbt.nhs.uk)