







Date of publication:09 April 2019Implementation:To be determined by each Service

# Change Notification UK National Blood Services No. 08 - 2019

# **Blood Safety Entry**

This change applies to the Whole Blood and Component Donor Selection Guidelines

Applies to England, Scotland and Wales only. Donors and staff in Northern Ireland see <u>Blood Safety Entry - N.I.</u>

Obligatory

Information must be provided so that individuals at risk do not donate. The reasons for donor self-exclusion must be understood.

#### 1. You must not donate if:

You think you need a test for HIV/AIDS, HTLV or hepatitis.

# 2. You must never donate if:

- a) You are HIV positive or receiving treatment for HIV.
- b) You are HTLV positive.
- c) You are a hepatitis B carrier.
- d) You are a hepatitis C carrier.
- e) You have ever injected, or been injected with, drugs; even a long time ago or only once. This includes bodybuilding drugs and injectable tanning agents. You may be able to give if a doctor prescribed the drugs. Please ask.

# 3. You must not donate for at least three months if:

You are working as a sex worker. You may be accepted for donation if it is longer than three months since you last received money or drugs for sex.

- 4. You must not donate for at least three months after sex (even if you used a condom or other protective) with:
- a) (If you are a man): another man.
- b) (If you are a woman): A man who has ever had oral or anal sex with another man, even if they used a condom or other protective. There are exceptions, so please ask.
- 5. You must not donate for at least three months after sex (even if you used a condom or other protective) with:

A partner who is, or you think may be:

- a) HIV or HTLV positive.
- b) A hepatitis B carrier.

# **\Continued**

Tel: 0114 358 4903 Fax: 0114 358 4494 Email: caroline.smith@nhsbt.nhs.uk









Tel: 0114 358 4903

Fax: 0114 358 4494

Email: caroline.smith@nhsbt.nhs.uk

- c) A hepatitis C carrier.
- d) A partner who has ever received money or drugs for sex.
- e) A partner who has ever injected, or been injected with, drugs: even a long time ago or only once. This includes bodybuilding drugs and injectable tanning agents. You may be able to give if a doctor prescribed the drugs. Please ask
- f) A partner who has, or you think may have been, sexually active, in parts of the world where HIV/AIDS is very common. This includes most countries in Africa. There are exceptions, so please ask.

#### 6. You must not donate for at least three months if:

- a) You have taken Pre-Exposure Prophlyaxis (PrEP) / Truvada® for prevention of HIV
- b) You have taken or been prescribed Post-Exposure Prophylaxis (PEP) for prevention of HIV.

If the underlying reason for taking PrEP or PEP warrants a longer deferral period, this should be applied.

#### See if relevant

Addiction and Drug Abuse Blood Safety Entry-N.I.

Hepatitis B Hepatitis C HIV

Homosexual and Bisexual Individuals

**HTLV** 

Infection - General

Non-consented Exposure to Human Bodily Fluids

Pre- or Post-Exposure Prophylaxis for HIV

### Additional Information

Following the publication of the SaBTO Donor Selection Criteria Report 2017 the deferral period for most risks in this category has been reduced from 12 to three months. In addition the permanent deferral of sex workers has been changed to three months.

This review considered advances in the sensitivity of testing procedures currently in use in the UK, the prevalence of transfusion transmissible infections and the current level of compliance with the 12 month exclusion for donors at higher risk of blood borne infections that has been in effect until now..

There may be exceptions for female partners of men who have ever had sex with men. They may be allowed to donate on the basis of an individual risk assessment.

The drugs used in both PrEP and PEP may interfere with the routine HIV screening tests carried out on all blood donations. For this reason, donors who have taken PrEP or PEP should not donate for three months, even if they have not had a sexual partner in a risk group.

'Blood Safety Leaflet Information' is available in the 'Document Library' of 'www.transfusionguidelines.org'.

#### **\Continued**









Information	This is a requirement of the Blood Safety and Quality Regulations 2005.
Reason for change	The addition of deferral periods for donors who have taken PrEP or PEP and a specific deferral of individuals on treatment for HIV.

Smaclernan

Dr Sheila MacLennan

<u>Professional Director</u> - Joint UKBTS Professional Advisory Committee

Tel: 0114 358 4903

Fax: 0114 358 4494