

Change Notification UK National Blood Services No. 13 - 2021

Immunoglobulin Therapy

This change applies to the Deceased Tissue Donor Selection Guidelines

Please amend the following sections of this entry:

Obligatory:	Must not donate if: a) Immunosuppressed. b) Donors with recovered immunodeficiency: Refer to a Designated Clinical Support Medical Officer.
Discretionary:	a) If the intravenous or subcutaneous human immunoglobulin was given before 1980, accept.
	b) Routine ante- and post- natal use of anti-D immunoglobulin, accept.c) If single dose prophylactic immunoglobulin has been given, accept.d) If treated with intravenous immunoglobulins after 1st January 1999: if
	underlying condition is not a contraindication, accept. Refer to designated clinical support officer if further advice required.
Additional Information:	Immunoglobulin used before 1980 is unlikely to be affected by vCJD. Single dose immunoglobulin is unlikely to pose a significant risk for transmitting vCJD.
	Since 1999, intravenous immunoglobulins prepared from UK donors have no longer been used, as a risk reduction measure for vCJD transmission.
Reason for Change:	The entry has been modified to comply with advice from the MSBTO committee of the DH. To permit donation from donors who have received intravenous immunoglobulin after 1 st January 1999, if the reason for treatment is not a contraindication.

\Continued









Update Information The advice reflects advice from the MSBTO committee of the DH.

Smachennan

 Dr Sheila MacLennan

 Professional Director

 - Joint UKBTS Professional Advisory Committee

 To Direct Dial : (0113) 820 8638

 Sheila.maclennan@nhsbt.nhs.uk