









## Change Notification UK National Blood Services No. 16 - 2012

## **Pregnancy**

Please modify the entries as follows.

Applies to the Tissue Donor Selection Guidelines Live Donors, Tissue Donor Selection Guidelines Deceased Donors and Bone Marrow and PBSC Donor Selection Guidelines

Obligatory Must not donate if:

Add an additional deferral criteria:

 c) It is less than 7 days from the last dose of methotrexate. (Live and Deceased DSG)

or

 e) It is less than 7 days from the last dose of methotrexate. (Bone Marrow and PBSC DSG)

Additional Information

Methotrexate is now increasingly used to medically treat ectopic pregnancy, to avoid surgery and protect the fallopian tube. A week is needed for any residual methotrexate to clear the system.

Add:

**Reason for Change** The addition of information about methotrexate.

## **Further Information**

The supporting paper, JPAC 12-15 Recommendations on a change to the Donor Selection Guidance for Pregnancy, leading to this Change Notification can be found in the Document Library/Supporting Papers of the JPAC website:

http://www.transfusionguidelines.org.uk/Index.aspx?Publication=DL&Section=12&pageid=7528

Dr Sheila MacLennan

Professional Director - Joint UKBTS/HPA Professional Advisory Committee

Direct Dial: (0113) 820 8638 Sheila.maclennan@nhsbt.nhs.uk

Maclena