

Issued by JPAC: 19 January 2021

Implementation: To be determined by each Service

## Change Notification UK National Blood Services No. 55 - 2020

### Tropical Virus

#### These changes apply to the Whole Blood and Components Donor Selection Guidelines

Please make the following changes:

<i>Definition</i>	To include Dengue Virus, Dengue Fever and Chikungunya Virus, also known as CHIKV, Zika Virus and Zika Virus Fever.  <b>Tropical Virus Endemic Areas:</b> are shown in the 'Geographical Disease Risk Index' (GDRI) as a Tropical Virus Risk.
<i>Obligatory</i>	<b>Must not donate if:</b> a) It is less than six months from a donor's return from a Tropical Virus Risk endemic area and the donor has been diagnosed with eChikungunya, eDengue or zZika virus infection whilst there or following their return to the UK. b) It is less than six months from a donor's return from a Tropical Virus Risk endemic area and the donor has either had a history of symptoms suggestive of eChikungunya, eDengue or zZika virus infection whilst there or following their return to the UK. c) In other cases it is less than four weeks from a donor's return from a Tropical Virus Risk endemic area. d) The donor has had sex in the last 28 days with someone who has had a confirmed Zika Virus infection in the <b>three 6</b> months preceding the sexual contact.
<i>Discretionary</i>	All donors may be accepted six months after their return from an affected area or resolution of symptoms. This may be reduced to four weeks, if they have had neither symptoms nor evidence of infection.
<i>See if Relevant</i>	Infection - General Malaria South American Trypanosomiasis The 'Geographical Disease Risk Index'

\Continued

<p><i>Additional Information</i></p>	<p>Chikungunya is an alpha virus that can cause a wide spectrum of disease. This may range from no or minimal symptoms to death. Most commonly it causes arthritis (typically in the knee, ankle and small joints of the extremities), high fever and a maculopapular rash.</p> <p>It is geographically widespread but since 2005 it has reached epidemic proportions in parts of India and islands in the Indian Ocean. It is known to be spread by blood in symptomatic cases and on theoretical grounds could be spread by transfusion and transplantation of tissues and organs from people with pre-symptomatic or asymptomatic disease. A number of visitors returning from endemic areas to the UK have been diagnosed with this infection.</p> <p>Dengue Virus is a flavivirus that typically gives rise to abrupt high fever with a range of accompanying symptoms. Dengue fever (DF) is the most common arthropod borne disease worldwide. Dengue is currently considered endemic in approximately 128 countries.</p> <p>Overall, 15-90% of cases may have an asymptomatic course of infection, but clinical presentation varies with age group. However there is a risk of change in disease presentation and potential for increased incidence of more severe disease in older age groups due to co-circulation of different dengue types and emergence of new types in endemic areas patterns.</p> <p>Zika virus is a flavivirus that is transmitted to humans through the bite of a carrier mosquito. Zika Virus can also be transmitted human to human through sexual contact. Zika infection is a rapid acute infection that in the majority of cases is asymptomatic or has very mild general symptoms. A small number of cases may have more apparent symptoms but hospitalisation is rare. Zika infection may be mistaken for Chikungunya or Dengue infections as the virus often co-circulate.</p> <p>The main vector for chikungunya virus, dengue virus and zika virus is <i>Aedes aegypti</i> (<i>Aedes albopictus</i> is another emerging vector), which is found worldwide between latitudes 35°N and 35°S. There is no epidemiologically important animal reservoir for Chikungunya, Dengue or Zika viruses. The main areas affected by all 3 viruses include the Caribbean, South and Central America, Mexico, Africa, the Pacific Islands, SE Asia, Indian sub-continent, Hawaii. Additionally Dengue fever has been reported in <del>Japan and</del> Australia <a href="#">and there have been outbreaks of Dengue and Chikungunya in Europe.</a></p> <p>As the problem can vary both in relation to geography and time of the year it is not possible to state areas from which donors need to be deferred or dates of disease activity. These are provided in the Geographical Disease Risk Index.</p> <p>Position statements are available in the JPAC Document Library.</p>
<p><i>Information</i></p>	<p>This entry is compliant with the Blood Safety and Quality Regulations 2005.</p>

\Continued

<p>Reason for change</p>	<p><del>Information about Zika virus transmission has been updated and deferral of donors with a history of sexual contact with someone known to have had a confirmed Zika virus infection in the 6 months prior to the sexual contact has been added.</del></p> <p>The deferral for sexual contact with an individual with confirmed Zika virus infection has been changed and now applies to partners with confirmed Zika infection in the last three months. Additional information now contains reference to European outbreaks.</p>
--------------------------	--

**Dr Sheila MacLennan**  
**Professional Director - Joint UKBTS Professional Advisory Committee**  
☎ Direct Dial: (0113) 820 8638    ✉ [sheila.maclennan@nhsbt.nhs.uk](mailto:sheila.maclennan@nhsbt.nhs.uk)