
IDENTIFICATION AND TREATMENT OF IRON DEFICIENCY ANAEMIA A guideline for use in primary & secondary care for elective surgery patients



KEY MESSAGES:

- Most anaemic patients have iron deficiency & the majority respond to oral or IV iron
- Patients referred for elective surgery must be screened for anaemia and their haemoglobin optimised before surgery, particularly where surgery is likely to involve moderate blood loss (>500 mls).
- For severe anaemia or symptomatic co-morbidities consider a minimal transfusion
- Refer to the appropriate secondary care specialism to investigate the underlying cause of anaemia e.g. GI, gynaecological, haematological, renal. *

EoE RTC V1 06.18

Examples of elective surgical procedures with the potential for blood loss

General surgery

Abdoperineal resection Anterior resection Bowel resection Cholecystectomy Gastrectomy Hemicolectomy Laparotomy Liver biopsy Oesophagectomy Oesophagogastrectomy Parathyroidectomy Splenectomy Thyroidectomy Vagotomy Whipples procedure

Orthopaedic surgery

Amputation Bilateral knee graft replacement Bone graft Elbow replacement Internal tibial fixation Knee replacement Osteotomy/bone biopsy Removal of nails Revision of knee replacement Revision of hip replacement Scoliosis correction Spinal decompressions Total hip replacement

Urology

Nephrectomy Percutaneous nephrolithotomy Prostatectomy Total cystectomy Trans- urethral resection prostate

Plastic surgery

Abdominoplasty Cosmetic breast reconstruction Head/neck reconstruction Mastectomy including radical

Vascular surgery

Above/below knee amputation Aortic aneurysm Axillo-femoral bypass Carotid endarterectomy Femoral endarterectomy Femoral-femoral cross over Femoral popliteal bypass Sympathectomy **Cardiac surgery** all referrals

Thoracic surgery

Exploratory thoracotomy Mediastinoscopy Open pleural biopsy Thoracotomy/pulmonary resection Tracheostomy Video assisted thoracic (VATS)

ENT

Block dissection neck/groin Bone graft Laryngectomy Osteotomy Parotid/submandibular gland Pharyngectomy

Gynaecology

Endometrial resection Hysterectomy Oophorectomy Vulvectomy

Obstetrics In collaboration with local antenatal department

Oral iron supplements dosage

Iron salt	Dose (daily) Minimum	Preparation	Ferrous iron content
Ferrous sulphate	200 mg	tablets	65mg
Ferrous gluconate	300 mg	tablets	35 mg
Ferrous fumarate	200 mg	tablets	65 mg
Sodium feredate	380mg/10 mls	elixir	55 mg