

Emily Graves Haem SpR

Please join in, take a guess, ask those 'stupid' questions...

Coming Up

- Why does blood need to be irradiated?
- Who needs irradiated products?
 - What happens if they get non-irradiated products?
- Whose responsibility is it anyway?

What condition are we trying to prevent when we irradiate?







Manifestations of TaGVHD



Rash



Severe diarrhoea (gut inflammation)



Jaundice (liver inflammation)

Fever





Death (in 100%)

Irradiation prevents TaGVHD by... stopping cell division.

It does not strengthen or mutate...

The Abomination and the Incredible Hulk, monsters created by accidental exposure to gamma irradiation. Not a side effect that is commonly seen in NHS practice.

Which products need irradiating?



Irradiation is performed at blood centres using either gamma or x-ray sources.

So, who is at risk?

Relevant to Adult Haem-Onc practice:

- Hodgkins Lymphoma
- Fludarabine recipients
 - □ CLL FCR, FMC, single agent
 - AML FLAG
 - □ NHL FC, single agent
- Other purine analogues/antagonists (lifelong)
 cladribine, pentostatin, clofarabine, bendamustine

(lifelong)

(lifelong)

(lifelong)

(lifelong?)

- Campath*
- Rabbit ATG**

^{*} Aplastic anaemia, CLL, some conditioning protocols

So, who is at risk?

Relevant to Adult Haem-Onc practice (cont'd)

- Autologous transplant recipients
 - 7 days prior to harvest, then from conditioning until 3 months post transplant
- Auto patients receiving Total Body Irradiation (for 6 months)
- Allogeneic transplant recipients
 - □ From conditioning until 6 months post transplant
 - □ OR: lymphocytes greater than 1
 - OR: GVHD prophylaxis or treatment discontinues; if have ongoing GVH then irradiation continues indefinitely
- Bone marrow donors (seven days prior to donation)

Who is at risk – rarer scenarios

- Donations from family members
- (HLA matched products)
- Intrauterine transfusions
- Neonatal exchange transfusions (if time allows)
- T lymphocyte deficiency syndromes (Di George etc)

What a long list – why not zap the lot?

- Red cells become leaky after irradiation potassium and free haemoglobin levels in the fluid increases.
- Therefore irradiated red cells can't be stored for as long as usual.
 - Must be less than 14 days old and then only stored for a maximum of 14 days, compared to usual red cell storage life of 35 days.
- And it's not cheap!

Whose responsibility is it anyway?



Who's responsibility is it anyway?

Communication is a big problem

Consultant/SpR

Pathologist

Pharmacist

Blood bank

Transplant centres

Prescribers

Nurse

Patient?





Information for patients needing irradiated blood

including important patient card and patient record stickers



The good news

- Universal leucodepletion was introduced in 1999.
- No cases of TaGVHD have been reported since 2001 (and only once since leucodepletion began)
- 686 cases of errors related to irradiated products reported to SHOT over this time.
- Leucodepletion may be enough to make blood is safe, but no-one is sure – so we can't forget irradiation!





A Few Cases

- Who made the mistake?
- What could be done differently?

Wife's knowledge overlooked by staff

A male undergoing chemotherapy for Hodgkin's disease was to receive his first transfusion. The request was made by a consultant haematologist, who failed to request irradiated components and did not supply the patient with an NHS card. When admitted to the haematology ward the following day, the wife questioned the need for irradiated components but was reassured by the nursing staff that this was not necessary.

Correspondence from Transplant Centre in medical records

A patient was referred back to his local hospital following a stem cell transplant. The Transplant Centre had written a discharge summary, which had included the requirement for the patient to receive both irradiated and CMV negative blood components. However, this correspondence was not available in the case notes at the time of the transfusion and was only passed on to the team from medical records several days later.

Patient on fludarabine receives non-irradiated red cells despite pharmacy protocol

A hospital transfusion laboratory receives a monthly report from the pharmacy of the patients who have been prescribed fludarabine. However, a locum haematologist prescribed fludarabine in the middle of the month and failed to give the patient an alert card or annotate the patient's case notes or inform the laboratory. Later that month the patient was transfused with 2 units of non-irradiated red cells, prior to the laboratory being aware through the monthly pharmacy report of the requirement.

THANK YOU!

Questions?

Sources, references

- BCSH Guideline 2010 (www.bcshguielines.com)
- Haemovigilance: www.shotuk.org
- The blood bank guy (<u>www.bbguy.org</u>)