



Original CXR



Splenic artery embolisation



Splenic artery embolisation




Blood Tests

Transfused 2 Units

Enoxaparin 90mg

[illegible]

DMSA Kidney scan at 4 months



Thank you

References

- Devon, Karen M., and Robin S. McLeod. "Pre and peri-operative erythropoietin for reducing allogeneic blood transfusions in colorectal cancer surgery." *Cochrane Database Syst Rev* 1 (2009).
- Madi-Jebara, Samia N., et al. "Postoperative intravenous iron used alone or in combination with low-dose erythropoietin is not effective for correction of anemia after cardiac surgery." *Journal of cardiothoracic and vascular anesthesia* 18.1 (2004): 59-63
- Cuenca, Jorge, et al. "Perioperative intravenous iron, with or without erythropoietin, plus restrictive transfusion protocol reduce the need for allogeneic blood after knee replacement surgery." *Transfusion* 46.7 (2006): 1112-1119.
- García-Erce, J. A., et al. "Perioperative stimulation of erythropoiesis with intravenous iron and erythropoietin reduces transfusion requirements in patients with hip fracture. A prospective observational study." *Vox sanguinis* 88.4 (2005): 235-243.
- Karkouti, Keyvan, et al. "Intravenous iron and recombinant erythropoietin for the treatment of postoperative anemia." *Canadian Journal of Anesthesia* 53.1 (2006): 11-19.
- Corwin HL, Parsonnet KC, Gettinger A. RBC transfusion in the ICU. Is there a reason? *Chest* 1995; 108: 767-71
- Smoller BR, Kruskall MS. Horowitz GL. Reducing adult phlebotomy blood loss with the use of pediatric sized blood collection tubes. *American Journal of Clinical Pathology* 1989; 91; 701-3