

## Guidelines for the Blood Transfusion Services

### 13.3: Pre-transfusion testing

<http://www.transfusionguidelines.org/red-book/chapter-13-patient-testing-red-cell-immunohaematology/13-3-pre-transfusion-testing>

### 13.3: Pre-transfusion testing

#### 13.3.1: Resolution of anomalous grouping

ABO grouping is the most important pre-transfusion serological test performed. Fully automated ABO and D grouping procedures have significantly improved the accuracy and security of results, and should be used wherever possible.

When anomalous ABO groups are encountered laboratory protocols should support investigation of the following findings.

Missing agglutinins in reverse grouping:

- obtain the patient's history, and review for information which may explain missing agglutinin (e.g. age, immunodeficiency, previous haemopoietic stem cell transplant)
- repeat the reverse group, increasing the sensitivity of the test, consider the use of tube techniques, lower incubation temperature, increased plasma:cell ratio and enzyme-treated red cells.

Unexpected additional reactions in the reverse group:

- investigate the presence of allo- or autoantibodies active at temperatures below 37°C
- consider repeating the reverse group at 37°C
- consider repeating the reverse group using cells negative for any identified alloantibody.

Unexpected reactions in the forward or D grouping, including positive diluent control:

- check for immunoglobulin coating of the patient's cells by performing a direct antiglobulin test (DAT)
- consider repeating tests using unpotentiated reagents in tube techniques
- consider techniques to remove or reduce immunoglobulin coating (e.g. warm wash or use 0.2M DTT to remove IgM) and repeat tests with appropriate controls.

Unexpectedly weak or mixed field reactions in forward or D group:

- obtain the patient's history, and review for information which may explain results (e.g. recent non-ABO identical transfusion, haemopoietic cell transplant)
- consider additional investigations which may include adsorption/elution, and flow cytometry

- panels of monoclonal anti-D reagents may be commercially available for the investigation of partial and weak D phenotypes.

Genotyping is useful in resolving grouping problems, particularly weak and partial D types (see section 15.2). Genotyping alone must not be used to determine the ABO group for use in selection of blood for transfusion. Where the patient ABO group cannot confidently be assigned by serology, group O (high-titre negative) blood must be selected.

### 13.3.2: Antibody identification

---

In all cases of the investigation of alloantibodies laboratories should focus on:

- secure identification of alloantibodies detected
- exclusion of additional specificities to those identified
- selection of blood for transfusion (Daniels *et al.* 2002)<sup>7</sup>

When antibodies which cannot be identified have been detected, laboratories should consider referral to the International Blood Group Reference Laboratory (IBGRL).

Antibody identification techniques and protocols are described in BCSH guidelines and should be adhered to. More complex problems encountered by RCI laboratories and not covered by BCSH are considered below.

#### 13.3.2.1: Complex antibody mixtures

When investigating complex antibody mixtures RCI laboratories should consider:

- extended phenotyping of the patient, e.g. C, c, D, E, e, K, M, N, S, s, P,  $\rho$ , Le<sup>a</sup>, Le<sup>b</sup>, Fy<sup>a</sup>, Fy<sup>b</sup>, Jk<sup>a</sup>, Jk<sup>b</sup>
- if this is not possible due to previous transfusion or IgG sensitisation, blood group genotyping should be considered
- extending the range of techniques and incubation temperature to identify component antibodies
- using cells matching the patient's phenotype/genotype to confirm the presence of multiple antibodies rather than an antibody to a high-frequency antigen
- careful use of alloadsorption/autoadsorption techniques to confirm the specificity of elements of the mixture

#### 13.3.2.2: Antibodies known as high-titre low-avidity (HTLA)

Antibodies traditionally known as HTLA include anti-Ch, -Rg, -Kn<sup>a</sup>, McC<sup>a</sup>, -Yk<sup>a</sup>, -Cs<sup>a</sup> and -Sl<sup>a</sup>. Typically HTLA antibodies present as reacting with most panel cells by indirect antiglobulin test (IAT) with variable strength, with or without similar patterns using enzyme-treated cells. Experienced operators can recognise characteristic agglutination by microscopic examination of tube IAT, which have been described as 'loose', 'stringy', 'fluffy', 'delicate' or 'gritty'. In investigating samples suspected to contain HTLA antibodies RCI laboratories should consider:

- Neutralising anti-Ch or -Rg specificities by incubating the patient's plasma with pooled group AB donor plasma before IAT is undertaken. Reactivity of these antibodies is usually abolished. A dilution control in which the patient's plasma is incubated with phosphate-buffered saline should be prepared

and tested in parallel with the neutralised plasma.

- The use of soluble recombinant CR1 proteins for inhibition studies
- The use of a panel of cells lacking HTLA antigens

### 13.3.2.3: Antibodies to high-frequency antigens (HFA)

Typically antibodies to HFA present with positive reactions of similar strength against all routine screen and identification panel cells. The most commonly encountered specificities include anti-k, –Lu<sup>b</sup>, –Kp<sup>b</sup>, –Vel, –Co<sup>a</sup>, –Yt<sup>a</sup>, –Fy3, –U and –In<sup>b</sup>. In investigating samples suspected to contain antibodies to HFA, RCI laboratories should consider:

- the ethnicity of the patient
- extended phenotyping as in section 13.3.2.1
- testing against a panel of rare reagent red cells
- typing the patient's red cells with antibodies to HFA. Where possible, CE-marked reagents must be used, otherwise results must be considered in context of the reliability of the reagent in use, supported by adequate controls.

### 13.3.2.4: Antibodies to low-frequency antigens (LFA)

Typically antibodies to LFA present with a negative antibody screen and are detected in crossmatch or investigation of HDFN. The most commonly encountered specificities include anti-Kp<sup>a</sup>, –Lu<sup>a</sup>, –Wr<sup>a</sup> and –Co<sup>b</sup>. In investigating samples suspected to contain antibodies to LFA, RCI laboratories should consider:

- testing the patient's plasma with a panel of red cells expressing LFA
- phenotyping the incompatible unit(s)/cells for LFA

## 13.3.3: Autoantibodies

---

Autoantibodies are frequently encountered in pre-transfusion testing, and may be the cause of autoimmune red cell destruction, or may be clinically benign. In either case autoantibodies may interfere with pre-transfusion testing, either due to coating of patient's cells with immunoglobulin, or as pan-reactive antibody in patient's plasma. In providing safe transfusion in the presence of autoantibodies, RCI laboratories may adopt the following strategies.

### 13.3.3.1: ABO and Rh grouping in the presence of autoantibodies

Most modern test systems support routine, accurate grouping of the majority of patients whose cells are coated with immunoglobulin and who give a positive DAT. Cases which are problematic may present with reaction patterns that cannot be assigned to an ABO group, weak additional reactions and positive reagent controls. Such cases should be investigated as in section 13.3.1.

Laboratories should make a clear documented assessment, based on the recommendations of reagent and test system suppliers, how to manage cases with anomalous ABO and D groups. This is particularly important when potentiated reagents are included in test systems.

### 13.3.3.2: Alloantibody detection and identification in the presence of autoantibodies

In dealing with cross-reacting autoantibodies, which complicate the detection and identification of underlying alloantibodies, RCI laboratories should consider:

- The characteristics of available, validated IAT in testing patient plasma-containing pan-reacting autoantibodies. Tube IAT may be less prone to interference by autoantibodies than column technologies.
- The use of the patient's own cells to adsorb autoantibody from the plasma, permitting detection and identification of alloantibodies.
- The use of cells from two or more selected donors to adsorb autoantibody. Typically these cells are enzyme treated to optimise removal of autoantibody. Alloantibodies to high frequency antigens are also likely to be removed by this technique.

#### **13.3.4: Management of patients with autoantibodies**

---

Consideration should be given to close matching of recipient and donor red cell types. This is to safeguard against the presence of alloantibodies undetected by tests on modified plasma, and to prevent further alloimmunisation. In patients who cannot be grouped by conventional serology, due to sensitisation of red cells or previous transfusion, genotyping offers a solution.

In patients with autoantibodies requiring regular transfusion, close matching of transfused red cells with the patient's own phenotype, to manage risk of transfusion reactions, may be used as a basis by scientists and clinicians to assess, and potentially reduce, the required frequency of testing. Such assessments should be fully documented and subject to planned review.

#### **13.3.5: Therapeutic monoclonal antibodies**

---

A number of conditions are treated using therapeutic monoclonal antibodies (TMAbs). These therapies have the potential to adversely interfere with serological investigations and compatibility testing in the blood bank, potentially causing unnecessary delays in providing blood components for transfusion. This may delay treatment of these patients, many of whom are transfusion dependent. Monoclonal antibody therapies may affect serological testing methods in a variety of ways, with the monoclonal antibody induced reactivity persisting for up to 6 months after the last treatment infusion.

The following testing protocols should be undertaken:

##### **13.3.5.1 Management of patients on therapeutic monoclonal antibodies**

Before TMAb therapy has started:

- Baseline ABO and D group (follow local policy for requirement of confirmatory sample rule for ABO and D group)
- Antibody screen, and antibody identification, if required
- Direct Antiglobulin Test (DAT)
- Extended phenotyping/genotyping for C, c, E, e, K, (k if K+), MNSs, Jk<sup>a</sup>, Jk<sup>b</sup>, Fy<sup>a</sup> and Fy<sup>b</sup> (genotyping to be used if the patient has been recently transfused, <3 months ago)

Once TMAb therapy has been commenced:

- ABO and D typing as per normal method (If the ABO group cannot be concluded, group O red cells may be required for transfusion)
- Antibody screening, and antibody identification if required, using a strategy to avoid the effect of the TMAb, e.g. reagent cells treated with 0.2M Dithiothreitol (DTT) used in patients on anti-CD38 TMAs
- Red cells should be matched for Rh (CcDEe) and K as well as for any alloantibodies
- In patients who have a strong panagglutinin, alloadsorption studies may allow satisfactory antibody detection / identification, alternatively use of a different serological technique, use of a soluble recombinant protein or use of an anti-human globulin reagent which lacks anti-IgG4 or use of rare phenotype red cells which lack the CD marker of interest may help with the resolution of the investigation.